

ORIGINAL ARTICLE

Use of the Ages and Stages Questionnaire and Bayley Scales of Infant Development-II in neurodevelopmental follow-up of extremely low birth weight infants

BJ Woodward¹, L-A Papile², JR Lowe³, VL Laadt³, ML Shaffer⁴, R Montman⁵ and KL Watterberg³

¹Department of Psychiatry and Behavioral Sciences, The Children's Hospital of Denver, Denver, CO, USA; ²Section of Neonatology, Department of Pediatrics, Baylor College of Medicine, Houston, TX, USA; ³Division of Neonatology, University of New Mexico Health Sciences Center, Albuquerque, NM, USA; ⁴Departments of Public Health Sciences and Pediatrics, Penn State College of Medicine, Hershey, PA, USA and ⁵General Clinical Research Center, University of New Mexico School of Medicine, Albuquerque, NM, USA

Objective: For infants born with extremely low birth weight (ELBW), we examined the (1) correlation between results on the Ages and Stages Questionnaire (ASQ) and the Bayley Scales of Infant Development-II (BSID-II) at 18 to 22 months corrected age; (2) degree to which earlier ASQ assessments predict later BSID-II results; (3) impact of ASQ use on follow-up study return rates.

Study Design: ASQ data were collected at 4, 8, 12 and 18 to 22 months corrected age. The BSID-II was completed at 18 to 22 months corrected age. ASQ and BSID-II 18 to 22 month sensitivity and specificity were examined. Ability of earlier ASQs to predict later BSID-II scores was examined through linear regression analyses.

Result: ASQ sensitivity and specificity at 18 to 22 months were 73 and 65%, respectively. Moderate correlation existed between earlier ASQ and later BSID-II results.

Conclusion: For extremely low birth weight infant assessment, the ASQ cannot substitute for the BSID-II, but seems to improve tracking success. *Journal of Perinatology* advance online publication, 10 February 2011; doi:10.1038/jp.2011.1

Keywords: bayley scales of infant development; ages and stages questionnaire; neurodevelopment; developmental assessment; developmental screening; NICU

Introduction

Long-term assessment of the outcomes of high-risk infants who received neonatal intensive care is an essential component of both understanding the implications of providing care to these patients and appropriately evaluating the results of interventions. The most

commonly used neurodevelopmental follow-up protocol for high-risk infants consists of a single, professionally administered neurodevelopmental evaluation at 18 to 22 months corrected age and includes administration of the Bayley Scales of Infant Development (BSID).^{1–3} This follow-up paradigm is expensive and difficult and frequently results in significant numbers of children lost to follow-up.^{4,5} If accurate information regarding developmental outcomes could be obtained from a parent-administered questionnaire, this could be both cost-effective and result in improved follow-up rates for high-risk infant follow-up studies. Studies indicate that parents can be reliable and valid sources of information regarding their child's current developmental status, when compared with professionally administered tests.^{6–8} Squires and Bricker were able to determine that most mothers, including those with risk factors such as low maternal education or a history of substance abuse, were able to use a parent-report screening tool and that their responses closely matched the results obtained by professionals using a standardized assessment measure.⁹

Previous comparisons of parent-administered developmental questionnaires with professionally administered neurodevelopmental evaluations of high-risk infants have yielded varying results.^{10–13} In this prospective follow-up study of extremely low birth weight infants, we asked to what degree the parent-administered Ages and Stages Questionnaire (ASQ) correlated with a professionally administered BSID-II assessment at 18 to 22 months corrected age; whether ASQs completed at 4, 8 and 12 months corrected age would predict performance on the BSID-II at 18 to 22 months corrected age; and whether use of the ASQ as a tool to maintain contact with families would improve follow-up rates.

Methods

Study population

Infants eligible for this study were 291 extremely low birth weight infants enrolled after parental consent into the PROPHET study

Correspondence: BJ Woodward, Department of Psychiatry and Behavioral Sciences, The Children's Hospital of Denver, Denver, CO 80045 USA.
E-mail: Woodward.Barbara@tchden.org
Received 25 May 2010; revised 22 December 2010; accepted 29 December 2010

and who survived to hospital discharge. The PROPHET Study was a multicenter, randomized, placebo-controlled trial of early and low-dose hydrocortisone to prevent bronchopulmonary dysplasia.¹⁴ Eligibility criteria for that study included birth weight between 500 and 999 g and the need for mechanical ventilation at 12 to 48 h of age.

Study procedures

Neurodevelopmental evaluation of infants enrolled in the PROPHET study included administration of the BSID-II and neurological examination by certified examiners at 18 to 22 months corrected age.³ Families were separately consented for the ancillary ASQ study and were asked to complete an ASQ when their child was 4, 8, 12 and 18 to 22 months corrected age. Approximately 2 weeks before an infant turning 4, 8, 12 and 18 to 22 months corrected age, an age-appropriate ASQ form was mailed to the home. Completed ASQs were either mailed back to the center (families were provided with stamped and addressed envelopes) or the research coordinator called the family and obtained the answers to the ASQ by phone. If the family had not completed the 18 to 22 month ASQ before the professional neurodevelopmental evaluation, the family was asked to complete the ASQ on site.

Developmental assessment tools

The BSID¹⁵ includes Mental (mental development index) and Psychomotor (psychomotor development index) Scales, as well as a Behavior Rating Scale. Raw scores on the BSID-II are converted to standardized scores with a mean of 100 and a s.d. of 15. For this study, a standard score of 70 or below, which is 2 s.ds below the mean, was considered a 'fail' on either the Mental or Psychomotor Scale of the BSID-II.

The ASQ: a parent-completed, child-monitoring system¹⁶ is a screening tool that uses parent observation to assess child development and behavior. Questionnaires are available at 2-month intervals from 4 to 24 months and at 30-, 33-, 36-, 48-, 54- and 60-months age points. The standardized assessment window begins 1 month before and ends 1 month following each questionnaire age point. Questions are written at a 4th to 6th grade reading level so that most parents can complete them independently. The questionnaires also can be used in an interview format for those who have difficulty with reading or language. The ASQ questionnaire at each age point contains six questions in each of five domains of development-communication, fine motor, gross motor, problem solving and personal social, for a total of 30 questions. Answer options for each question include 'yes', 'sometimes' or 'not yet'. A 'yes' response receives 10 points, 'sometimes' receives 5 points and 'not yet' receives 0 points. Each of the five domains is scored separately. These separate scores are not added together to obtain a composite score. A score of 2 s.ds or more below the mean in any one of the domains is considered a 'fail' on the ASQ.¹⁶

Statistical analysis

To examine the relationship between the 18 to 22 month ASQ and the BSID-II, a 2X2 contingency table was constructed where the rows and columns were determined by a pass or fail of the ASQ and BSID-II, respectively. From this table, the test characteristics of the ASQ were calculated, including the sensitivity, specificity and positive and negative predictive values. The ability of the ASQ to predict performance on the Bayley Mental Scale and the Bayley Psychomotor Scale was examined using linear regression analysis without adjustment for other factors. The Spearman's rank correlation coefficient was used to measure the degree of association between the percentage of subjects with two or more ASQs completed and the follow-up rate. Analyses were conducted using SAS Version 9 (SAS Institute, Cary, NC, USA).

Results

Of the 291 infants who survived to discharge, 250 completed the BSID-II at 18 to 22 months corrected age. Of the 250 children who completed the BSID-II, 219 also completed at least one of the possible four ASQ questionnaires. Table 1 shows the eight unique ASQ patterns of return across the four age points. Of the 228 children, a total of 78 ASQs were completed at 4 months, 95 at 8 months, 105 at 12 months and 228 at 18 to 22 months. As illustrated, nine of the children with one or more ASQs did not complete the BSID-II evaluation. Characteristics for the 228 children in this study included: mean birth weight—738.5 g

Table 1 Patterns of return of the ASQs

Pattern of ASQ Return	ASQ Completion and Return Age Points					
	ASQs completed at 4 months.	ASQ completed at 8 months.	ASQ completed at 12 months.	ASQ completed at 18 to 22 mos.	Also the BSID-II Exam	Not the BSID-II Exam
	A	0	0	0	65	1
B	0	0	27	27	3	
C	0	15	0	15	0	
D	21	0	0	21	3	
E	0	34	34	34	1	
F	11	0	11	11	1	
G	13	13	0	13	0	
H	33	33	33	33	0	
Totals	78	95	105	219	9	

Abbreviations: ASQs, Ages and Stages Questionnaires; BSID-II, Bayley Scales of Infant Development-II.
n = 228 children.

Eight patterns of ASQ return are illustrated across 4 age points: 4, 8, 12 and 18 to 22 months. The 18 to 22 month column indicates whether the BSID-II exam was completed.

Table 2 Sensitivity/specificity of the 18 to 22 month ASQ and BSID-II

	BSID-II		Total
	Pass ^a	Fail ^b	
<i>(a)</i>			
ASQ			
Pass ^a	71	16	87
Fail ^b	39	43	82
Total	110	59	169
<i>(b)</i>			
	BSID-II		Total
	Pass ^a	Fail ^b	
ASQ			
Pass ^c	35	3	38
Fail ^d	75	56	131
Total	110	59	169
<i>(c)</i>			
	BSID-II		Total
	Pass ^e	Fail ^f	
ASQ			
Pass ^a	47	40	87
Fail ^b	16	67	83
Total	63	107	170

Abbreviations: ASQs, Ages and Stages Questionnaires; BSID-II, Bayley Scales of Infant Development II; NPV, negative predictive value; PPV, positive predictive value.

(a): Sensitivity: 73% (PPV = 52%); Specificity: 65% (NPV = 82%).

(b): Sensitivity: 95% (PPV = 43%); Specificity: 32% (NPV = 92%).

(c): Sensitivity: 63% (PPV = 81%); Specificity: 75% (NPV = 54%).

^a ≤ 2 s.d. below the mean.

^b > 2 s.d. below the mean.

^c ≤ 1 s.d. below the mean.

^d > 1 s.d. below the mean.

^e ≤ 1 s.d. below the mean.

^f > 1 s.d. below the mean.

(500 to 997 g.); mean gestational age—25.4 weeks (23.0 to 31.0 weeks); gender—males 121 (53.1%), females 107 (46.9%); ethnicity—Non-Hispanic white 117 (51.3%), Black 73 (32.0%), Hispanic 27 (11.8%), other 11 (4.8%); median total household income was \$30 000 to \$40 000.

Relationship between the BSID-II and ASQ at 18 to 22 months

Table 2a presents the test characteristics of the ASQ, including the sensitivity, specificity, and positive and negative predictive values. When using cutoff points of >2 s.d. below the mean on either the Mental Development Index or Psychomotor Development Index of

the BSID-II as a ‘fail’ and a score of >2 s.d. below the mean on any of the domains of the ASQ as a ‘fail’, the ASQ had a sensitivity of 73% and a specificity of 65%. The positive predictive value was 52% and the negative predictive value was 82%. In all, 16 of 59 children who scored poorly (>2 s.d. on either the Mental or Psychomotor Scale) on the BSID-II would have been missed by the ASQ, whereas 39 of 110 children would have been incorrectly identified as having a developmental delay. Thus, the clinical epidemiological characteristics (sensitivity and specificity) of the ASQ in relationship to the BSID-II did not reach suggested levels of 80% (see ref. 17).

We evaluated the effect of using a ‘fail’ cut-off on the ASQ of >1 s.d. below the mean, rather than >2 s.d. Table 2b shows the corresponding shifts in the sensitivity, specificity, positive and negative predictive values. The shift from 2 s.d. to 1 s.d. below the mean resulted in improved sensitivity with the ASQ only ‘missing’ three children with a developmental delay. On the other hand, 75 children would have been incorrectly considered as suspect for a developmental delay but were found to be above the BSID-II cut off of 2 s.d. below the mean, thus decreasing the specificity to 32%.

We also examined the effect on sensitivity and specificity rates by shifting the s.d. of the BSID-II to >1 s.d. below the mean, while keeping the ASQ at >2 s.d. We found that the sensitivity rate dropped to 63%, whereas the specificity improved slightly to 75% (see Table 2c).

Early ASQ prediction of 18–22-month BSID-II

As illustrated in Table 1, a total of 154 children had at least one completed ASQ at age 4, 8 and/or 12 months, plus completed the BSID-II at 18 to 22 months corrected age. Excluded from this sub-sample were the 65 children who had no ASQ results before the 18 to 22 month BSID-II evaluation and nine children who had ASQs completed but were not brought in for the BSID-II evaluation. As illustrated in Table 1, the number of ASQs completed per child over the three age points of 4, 8 and 12 months was variable; therefore, the number of ASQs available for calculations at each age point is different.

Tables 3 and 4 present the regression results as coefficients of multiple correlation using combinations of the ASQ at 4, 8 and 12 months to predict performance on the 18 to 22 month Bayley Mental Scale (Table 3) and Psychomotor Scale (Table 4). Overall predictability was mild to moderate for each of the scales, indicating that the majority of the variability in the BSID-II still needs to be explained after accounting for ASQ performance. The combination of the 4 and 12 month ASQ data points was the strongest predictor of the 18 to 22 month BSID-II Mental Scale ($R^2 = 0.23$; $R = 0.48$) and Psychomotor Scale ($R^2 = 0.32$; $R = 0.57$).

ASQ use as a population tracking and retention mechanism

The rates of return for the 18 to 22 month follow-up visit were compared with numbers of completed ASQs per participating

Table 3 Coefficients of multiple correlation between ASQs and BSID-II MDI

Age at ASQ (months)	n	R	BSID-II Mean	P-value
4	73	0.41	80.55	0.002
8	93	0.45	81.72	<0.001
12	103	0.45	81.76	<0.001
4+8	45	0.29	83.09	0.050
4+12	42	0.48	84.62	0.001
8+12	67	0.44	83.33	<0.001
4+8+12	33	0.47	85.70	0.006

Abbreviations: ASQs, Ages and Stages Questionnaires; BSID-II, Bayley Scales of Infant Development II; MDI, Mental development index.

research center. Using a Spearman's correlation coefficient, a significant positive correlation ($r_s = 0.71$) was found when two or more ASQs were completed per child before the 18- to 22-month evaluation as shown in Figure 1.

Discussion

Although the ASQ holds promise for supporting the efforts of measuring the development of young children born with extremely low birth weight, the clinical epidemiological characteristics (sensitivity and specificity) of the ASQ in relationship to the BSID-II did not reach suggested levels of 80% (see ref. 17). Although the values came close to meeting suggested guidelines for accurate classification of children, the number of children who would be correctly identified as not having a developmental delay (specificity) is not strong enough to assert that the ASQ provides accurate classification when compared with the BSID-II. Specificity values declined even further when the cutoff score for failure on the ASQ was shifted from 2 s.d. to 1 s.d.

Skellern, *et al.*¹³ reported overall sensitivity and specificity rates for the ASQ of 90 and 77%, respectively. Their study used an ASQ cutoff score of >2 s.d., but adjusted the cutoff score of the evaluation tools (BSID-II, Griffith Mental Development Scales, and the McCarthy Scales of Children's Abilities General Cognitive Intelligence Scale) to >1 s.d. When we shifted to a combination of ASQ cutoff scores of >2 s.d. and cutoff scores of >1 s.d. for the BSID-II, sensitivity dropped to 63%, whereas the specificity improved toward acceptable levels (75%). It is interesting to note that when Skellern *et al.* used either the Griffith Mental Development Scales or the McCarthy scales of Children's Abilities General Cognitive Intelligence Scale, they achieved much higher sensitivity and specificity values for the ASQ than they did when using the BSID-II as the index evaluation tool. A number of studies have attempted to determine whether scores obtained during the first 2 years of life, using professionally administered developmental measures such as the BSID-II, can predict later outcomes. Most of these studies reveal weak correlations.^{18–21}

Table 4 Coefficients of multiple correlation between ASQs and BSID-II PDI

Age at ASQ (months)	n	R	BSID-II Mean	P-value
4	75	0.46	85.75	<0.001
8	93	0.36	85.97	0.002
12	104	0.44	86.21	<0.001
4+8	45	0.40	88.84	0.006
4+12	43	0.57	89.95	<0.001
8+12	67	0.41	86.91	<0.001
4+8+12	33	0.44	90.42	0.011

Abbreviations: ASQs, Ages and Stages Questionnaires; BSID-II, Bayley Scales of Infant Development II; PDI, Psychomotor Development Index.




Figure 1 Per center comparison between the percentage of children who had two or more Ages and Stages Questionnaires (ASQs) with the percentage of Bayley Scales of Infant Development II (BSID-II) evaluations completed ($r_s = 0.71$). Each data point represents one research center.

For this study, the ASQs completed at 4, 8 and 12 months had mild to moderate correlations with the Mental Development Index and Psychomotor Development Index of the BSID-II administered at the 18 to 22 month corrected age point. It is interesting to note that correlations with the Psychomotor Development Index improved when using either the 4- or 12-month ASQ results, or when the 4- and 12-month ASQ scores were added in combination.

The use of the ASQ seemed to support efforts to retain infants in the PROPHET study until the time of the 18 to 22 month professionally administered developmental evaluation. A significant positive correlation ($r_s = 0.71$) was found between a family's completion of two or more ASQs and their return for the professional evaluation. The possibility exists, however, that families who are motivated to return to a center for an evaluation of their child's development, also would be more likely to complete and return ASQ questionnaires. It also should be noted that more frequent family contact in any form improved follow-up rates. One urban center had a high follow-up rate with few ASQs completed.

Their protocol was to bring children in to the follow up clinic every 3 months. Although frequent returns to the hospital/clinic may be physically and financially possible in more urban areas, this might not be feasible in more rural settings.

Most of the research coordinators indicated that they liked using the ASQ because it gave them a reason for contacting the families. They related that families who completed the ASQs appreciated the knowledge gained about their child's developmental progress. Several coordinators indicated that information gathered through the ASQ process helped families to better understand their child's developmental status. In this way developmental findings determined at the study evaluation were not unexpected and thus were less traumatic for the family.

One limitation of this study was the relatively small number of ASQs at each of the earlier age points. This fact might have influenced the results regarding the ASQ's ability to predict BSID-II scores at 18–22 months. These smaller numbers also did not allow for examination of any possible influences related to sociodemographical factors.

In addition, it should be noted that since the completion of this study, the ASQ-3 (see ref. 22) and the BSID-III (see ref. 23) have been published. The ASQ-3 has additional age intervals, cutoff scores have been revised, and the scoring sheet displays the cutoff point for 1 s.d. below the mean. The BSID-III contains five subtests including motor, language, cognitive, social-emotional and adaptive behavior, which may allow for more direct comparisons between the domains of the ASQ-3 and the BSID-III in future studies.

Another relevant comparison worthy of consideration is that between the BSID-III and the BSID-II (see ref. 24), where significant differences have been shown. Mean composite scores on the BSID-III are reported to be ~7 points higher than mean index scores on the BSID-II (see ref. 24). Depending upon one's view of the relative strengths of each tool, either the BSID-II overestimates delays or as Anderson *et al.*²⁵ indicate, the BSID-III 'seriously underestimates' developmental delay. It seems that more work needs to be done to discover which of the available assessment tools, the BSID-II, BSID-III, ASQ or others, best document current developmental status and in addition is the best predictor of child outcomes.

In summary, the results of this study indicate that the ASQ does not have strong enough clinical epidemiological properties to support replacing the BSID-II for neonatal intensive care unit (NICU) follow-up studies. More positively the results of the ASQ may give researchers some indication of the child's development if the family does not return for the professionally administered evaluation. The ASQ also helps families to feel included in the process of monitoring their child's development and provides them with valuable developmental information. It is possible that the periodicity of the ASQ process may encourage families to continue their connection with the research center and thus may be a useful

adjunct to prospective neurodevelopmental research outcome studies as an effective tracking and population-retention mechanism.

Accurate determination of developmental outcomes of newborns cared for in the neonatal intensive care unit is critical to informing ongoing care. However, the question must be posed as to whether the current paradigm of a one-time professionally administered evaluation, before 2 years of age, is yielding accurate information about developmental outcomes. As previously noted, developmental evaluations performed in the early years have not been found to be predictive of later outcomes.^{18–21} The question needs to be raised as to whether a periodic parent-administered measure, such as the ASQ, may yield better predictive information. Studies contrasting early developmental findings gathered through the use of professionally administered evaluation tests with results from parent-completed measures would need to be conducted to determine each method's ability to predict long-term developmental outcomes.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

This study was conducted at the University of New Mexico, State University of New York at Buffalo, University of Colorado, Johns Hopkins University, St Joseph Regional Medical Center of Milwaukee, Children's Hospitals and Clinics of Minneapolis and St Paul, MN, Tufts University, University of Pennsylvania, and Virginia Commonwealth University. The study was supported by a grant from the National Institute of Child Health and Human Development (R01—HD38540) and grants from the General Clinical Research Centers Programs at the University of New Mexico (M01 RR00054), Tufts-New England Medical Center (5M01 RR00997), and the University of Colorado (M01-RR00069).

References

- 1 Vohr BR, Wright LL, Dusick AM, Mele L, Verter J, Steichen JJ *et al.* Neurodevelopmental and functional outcomes of extremely low birth weight infants in national institute of child health and human development neonatal research network, 1993-1994. *Pediatrics* 2000; **105**: 1216–1225.
- 2 Ohls RK, Ehrenkranz RA, Das A, Dusick AM, Yolton K, Romano E *et al.* Neurodevelopmental outcome and growth at 18-22 months corrected age in extremely low birth weight infants treated with early erythropoietin and iron. *Pediatrics* 2004; **114**: 1287–1291.
- 3 Watterberg KL, Shaffer ML, Mishefske MJ, Leach CL, Mammel MC, Couser RJ *et al.* Growth and neurodevelopmental outcomes after early low-dose hydrocortisone treatment in extremely low birth weight infants. *Pediatrics* 2007; **120**: 40–48.
- 4 Hille ET, Elbertse L, Grovenhorst JB, Brand R, Verloove-Vanhorick SP. Nonresponse bias in a follow-up study of 19-year-old adolescents born as preterm infants. *Pediatrics* 2005; **116**(5): e662–e666.
- 5 Aylward GP. Methodological issues in outcome studies of at risk infants. *J Pediatr Psychol* 2002; **27**: 37–45.

- 6 Glascoe FP, Dworkin PH. The role of parents in the detection of developmental and behavioral problems. *Pediatrics* 1995; **95**: 829–836.
- 7 Glascoe FP. Parents' concerns about children's development: prescreening technique or screening test? *Pediatrics* 1997; **99**: 522–528.
- 8 Diamond KE, Squires JK. The role of parental report in the screening and assessment of young children. *J Early Interv* 1993; **17**: 107–115.
- 9 Squires J, Bricker D. Impact of completing infant developmental questionnaires on at-risk mothers. *J Early Interv* 1991; **15**(2): 162–172.
- 10 Kim M, O'Connor K, McLean J, Robson A, Chance G. Do parents and professionals agree on the developmental status of high-risk infants? *Pediatrics* 1996; **97**: 676–681.
- 11 Klammer A, Lando A, Pinborg A, Greisen G. Ages and Stages Questionnaire used to measure cognitive deficit in children born extremely preterm. *Acta Paediatr* 2005; **94**: 1327–1329.
- 12 Plomgaard AM, Hansen BM, Greisen G. Measuring developmental deficit in children born at gestational age less than 26 weeks using a parent-completed developmental questionnaire. *Acta Paediatr* 2006; **95**: 1488–1494.
- 13 Skellern C, Rogers Y, O'Callaghan MJ. A parent-completed developmental questionnaire: follow up of ex-premature infants. *J Paediatr Child Health* 2001; **37**: 125–129.
- 14 Watterberg KL, Gerdes JS, Cole CH, Aucott SW, Thilo EH, Mammel MC *et al*. Prophylaxis of early adrenal insufficiency to prevent bronchopulmonary dysplasia: a multicenter trial. *Pediatrics* 2004; **114**: 1649–1657.
- 15 Bayley N. *Bayley Scales of Infant Development*, 2nd edn. The Psychological Corporation: San Antonio, TX, 1993.
- 16 Bricker D, Squires J. *Ages and Stages Questionnaire (ASQ): A Parent-Completed, Child Monitoring System*, 2nd edn. Brookes Publishing: Baltimore, MD, 1999.
- 17 Glascoe FP. Standards for screening test construction. 2004. Available at: www.dbpeds.org. Accessed 05/04/09.
- 18 Aylward GP, Gustafson N, Verhulst SJ, Colliver JA. Consistency in the diagnosis of cognitive, motor, and neurologic function over the first three years. *J Pediatr Psychol* 1987; **12**: 77–98.
- 19 McGrath E, Wypij D, Rappaport LA, Newburger JW, Bellinger DC. Prediction of IQ and achievement at age 8 years from neurodevelopmental status at age 1 year in children with D-Transposition of the great arteries. *Pediatrics* 2004; **114**: e572–e576.
- 20 Hack M, Taylor HG, Drotar D, Schluchter M, Cartar L, Wilson-Costello D *et al*. Poor predictive validity of the Bayley Scales of Infant Development for cognitive function of extremely low birth weight children at school age. *Pediatrics* 2005; **116**: 333–341.
- 21 Roberts G, Anderson PJ, Doyle LW. The stability of the diagnosis of developmental disability between ages 2 and 8 in a geographic cohort of very preterm children born in 1997. *Arch Dis Child* 2010; **95**(10): 786–790.
- 22 Squires J, Bricker D. *Ages and Stages Questionnaire*, 3rd edn. (ASQ-3TM) Brookes Publishing: Baltimore, MD, 2009.
- 23 Bayley N. *Bayley Scales of Infant Development*, 3rd edn. PsychCorp, Harcourt Assessment, Inc.: San Antonio, TX, 2006.
- 24 Bayley N. *Bayley Scales of Infant Development*, 3rd edn., Technical Manual PsychCorp, Harcourt Assessment, Inc.: San Antonio, TX, 2006.
- 25 Anderson PJ, De Luca CR, Hutchinson E, Roberts G, Doyle LW. Underestimation of developmental delay by the new Bayley-III Scale. *Arch Pediatr Adolesc Med* 2010; **164**(4).